

DRAFT

**INTERACTION BETWEEN SMOKING AND
OCCUPATIONAL EXPOSURES**

National Institute for Occupational
Safety and Health

Authors

Dr. Jean G. French
Dr. Harvey P. Stien
Dr. William J. McKay
Dr. Bruce E. Albright
Dr. George E. Casey
Ms. C. Ilana Howarth

1005058068

INTERACTION BETWEEN SMOKING AND OCCUPATIONAL EXPOSURES

2

3

4

5

INTRODUCTION

7

There is increasing evidence of interactions between the smoking of tobacco and workplace exposures in the development of certain disease states. Most authorities believe that smoking contributes substantially to the occurrence of a spectrum of diseases including chronic respiratory disease, cardiovascular disease, and specific types of cancer, but it has not been generally recognized that occupational exposure to certain physical and chemical agents also contribute to the development of the same disease states. Some of the effects attributed to smoking may reflect an interaction between smoking and occupational exposure. This can not be quantified at the present time, but at least six different ways have been identified in which smoking may act with physical and chemical agents found in the workplace. These actions are not mutually exclusive and several may prevail for any given agent.

Six ways in which smoking may act with physical and chemical agents to produce adverse health effects are:

1. Tobacco products may serve as vectors by becoming contaminated with toxic agents found in the workplace, thus facilitating entry of the agent by inhalation, ingestion, and/or skin absorption.

10050580639

2. Workplace chemicals may be transformed into more harmful agents by smoking. 31 32
3. Certain compounds in tobacco products and/or in the smoke may be the same as toxic agents found in the workplace, thus increasing exposure to the agent. 34 35 36
4. Smoking may contribute to an effect comparable to that resulting from exposure to toxic agents found in the workplace, thus causing an additive biological effect. 39 40
5. Smoking may act synergistically with toxic agents found in the workplace to cause a more profound effect than that resulting from the agent and smoking added together. 43 44 45
6. Smoking may contribute to accidents in the workplace. 47

Exposure to multiple physical and chemical agents in the workplace can compound these various types of actions. 50

1005058070

ILLUSTRATIVE EXAMPLES OF DIFFERENT TYPES OF ACTIONS BETWEEN SMOKING AND 57
OCCUPATIONAL EXPOSURES

Tobacco products may serve as vectors by becoming contaminated with 59
toxic agents found in the workplace, thus facilitating entry of 60
the agent by inhalation, ingestion, and/or skin absorption. 61

Workplace chemicals may be transformed into more harmful agents by 65
smoking. 66

Investigations of outbreaks of polymer fume fever provide clear 68
illustrations of tobacco products serving as vectors for workplace 69
chemicals. In addition, these case studies demonstrate that workplace 70
chemicals can be transformed into more toxic agents by tobacco smoking. 72

Polymer fume fever is a disease with influenza-like symptoms caused by 75
inhalation of fumes from heated polytetrafluoroethylene (1). Typical 76
symptoms include chest discomfort, fever, leukocytosis, headache, 77
chills, muscular aches, and weakness. Since the symptoms are so similar 78
to influenza, polymer fume fever is difficult to diagnose. Workers who 79
continue to smoke may experience continuing reexposure and persisting 80
symptomatology. Although complete recovery has usually been reported to 81
occur within 12-48 hours after exposure is terminated, an autopsy report 82
attributes permanent lung damage to repeated episodes of polymer fume 83
fever (2). Pulmonary edema following exposure to heated 84
polytetrafluoroethylene has also been reported (3,4). Polymer fume 86
fever was first described in the literature in 1951 (5) as a result of

1005058071

two workers being exposed to polytetrafluoroethylene heated to 450- 87
500° C. The particular decomposition products(s) responsible for polymer 88
fume fever have not yet been identified, but temperatures in excess of 89
315° C have been sufficient to cause symptoms. The temperature of the 90
combustion zone of cigarettes is approximately 875° C (6). 91
Numerous outbreaks of polymer fume fever among smokers have been 94
attributed to the decomposition of workplace polytetrafluoroethylene by 95
lit cigarettes and inhalation of the harmful decomposition products with 96
cigarette smoke. One report (7) describes aviation employees whose work 97
involved contact with door seals that had been sprayed with an 98
unspecified fluorocarbon polymer. In one case, a worker smoking during
a break realized by the taste of his cigarette that it had become 99
contaminated. Although the worker extinguished the cigarette, he 100
experienced shivering and chills, which lasted approximately six hours, 101
beginning a half hour after this incident. Another illustrative report 102
(8) describes outbreaks of polymer fume fever among smoking workers 103
whose hands were contaminated with polytetrafluoroethylene used as a 104
mold release agent. There was no recurrence of symptoms after smoking 105
at the plant was prohibited. An outbreak of polymer fume fever among 106
workers using liquid fluorocarbon polymer in the production of imitation 107
crushed velver was likewise attributed to decomposition of fluorocarbon 108
polymer by lit cigarettes (9). Processing temperatures at this plant 109
were too low to pyrolyze the polymer. The seven affected workers were 110
all cigarette smokers whereas most of the workers without symptoms were 111
non-smokers. After work practices were changed to prohibit smoking in 112
the work area and to require hand washing before smoking, no further 113
symptoms at this facility were reported. Other outbreaks of polymer 114

1005058072

fume fever attributed to cigarette smoking have also been reported
(10,11,12,13,14).

The effects of smoking cigarettes contaminated with known amounts of 117
tetrafluoroethylene polymer have been studied with the assistance of 118
human volunteers. (15) Nine out of ten subjects were reported to 119
exhibit typical polymer fume fever symptoms after each had smoked just 119
one cigarette contaminated with 0.40 mg tetrafluoroethylene polymer. 120
Onset of symptoms ranged from 1 to 3.5 hours after smoking; recovery 121
time averaged nine hours.

Additional research is clearly warranted to identify other workplace 123
chemicals which are transformed into more toxic agents by tobacco 124
smoking.

With respect to tobacco products serving as vectors, the National 127
Institute for Occupational Safety and Health (NIOSH) has thus far 128
identified the following agents as potential candidates for 129
contamination of tobacco and tobacco products. 130

132

1005058073

<u>Agent</u>	<u>Major Health Effects</u>	133
		134
Formaldehyde (16)	Respiratory irritant, dermatitis	135
Boron Trifluoride (17)	Respiratory irritant, joint disease	136
Organotin (18)	Respiratory irritant	137
Methyl Parathion (19)	Reduced erythrocyte cholinesterase activity	138 139
Dinitro-ortho-Creosol (20)	Kidney damage, peripheral neuritis, CNS disturbances.	140 141
Carbaryl (21)	Inhibition of acetylcholinesterase	142
Inorganic Fluorides (22)	Fluoride osteosclerosis	143
Inorganic Mercury (23)	CNS disturbances, kidney damage, peripheral neuritis	144 145
Lead (24,25)	Nervous system toxin, renal toxin, changes in hematopoietic system	146 147 148 149

1005058074

Certain compounds in tobacco products and/or smoke may be the same as 152
toxic agents found in the workplace, thus increasing exposure to the agent. 153

Hydrogen Cyanide 156

Hydrogen cyanide has been found in cigarette smoke at concentrations as 159
high as 1,600 ppm (26). In 1973 Pettegrew and Fell (27) found the 160
plasma thiocyanate (a metabolite of cyanide) levels of smokers
significantly elevated as compared to those in non-smokers. In 1973 161
Radojicic (28) reported a study of 43 workers in the electroplating 162
division of an electronics firm in Nes, Yugoslavia. He found that the 164
majority of workers exposed to cyanide complained of fatigue, headache, 165
asthenia, tremors of the hands and feet, and pain and nausea. The 166
urinary thiocyanate concentrations of the exposed group of workers were 167
higher at the end of the work shift than before exposure at work.

Urinary thiocyanate concentrations were significantly higher among 168
exposed smokers than unexposed smoking controls, significantly higher 169
among exposed non-smokers than unexposed non-smokers, and significantly 170
higher among exposed smokers than among exposed non-smokers. These 171
findings demonstrate that smoking and occupational exposure can each 172
contribute to a workers' total exposure to and intake of cyanide. 173

Adverse effects from cyanide may occur from sublethal fatal doses. 175

Hydrogen cyanide and cyanide salts inhibit cytochrome oxidase. Cyanide 178

1005058075

can form complexes with heavy metal ions. Formations of these complexes 179
in the body can rapidly cause disturbances in enzyme systems in which 180
heavy metals act as cofactors either alone or as part of organic 181
molecules (29,30,31). Thiocyanate itself has toxic effects, especially 182
inhibition of uptake of inorganic iodide into the thyroid gland for 183
incorporation into thyroxin (32). The National Institute for 184
Occupational Safety and Health has estimated that over twenty thousand 185
workers in seventy-five different occupational groups have potential 186
occupational exposure to cyanide (33).

188

189

Carbon Monoxide (CO) 192

Cigarette smoking causes increased exposure to CO. A CO concentration 195
of 4 percent (40,000 ppm) in cigarette smoke generates an alveolar CO 196
concentration of 0.04 to 0.05 percent (400 to 500 ppm) which produces a 197
carboxyhemoglobin (COHb) concentration of 3 to 10 percent (34,35,36).

Goldsmith (21) estimated that the cigarette smoker is exposed to 475 ppm 198
CO for approximately six minutes per cigarette. 199

In a study on the COHb levels in British steelworkers, Jones and Walters 202
(38) found a 4.9 percent end of shift COHb saturation in non-smoking 203
blast furnace workers compared to 1.5 percent saturation in non-smoking 204
unexposed controls. For heavy cigarette smokers, the levels were 7.4 205

1005058076

percent for blast furnace workers and 4.0 percent for smoking unexposed 206
controls. The carboxyhemoglobin levels of blast furnace workers who
smoked were in a critical range. Studies by Knelson (39) and Hovarth 207
(40) have shown that levels of COHb in excess of 5 percent can cause 208
cardiovascular alterations which are dangerous for persons with coronary 210
heart disease.

Potential occupational exposure to carbon monoxide is great (41). Since 214
a significant number of workers with coronary heart disease do smoke,
additional occupational exposure to carbon monoxide may increase 215
-cardiovascular morbidity and mortality. 216

Methylene Chloride 219

Methylene Chloride is metabolized to CO in the body (42). 221
Carboxyhemoglobin levels in blood increase with increasing environmental 222
concentrations of methylene chloride as well as with increasing physical 223
activity at the time of exposure (43,44). Maximum carboxyhemoglobin 224
levels occur 3 to 4 hours after exposure is discontinued. 225

Mean methylene chloride concentrations of 778 ppm over a three hour 227
exposure period produced a maximum carboxyhemoglobin level of 9.1% four 228
hours after exposure was discontinued. Twenty hours after this exposure 229

1005058077

the carboxyhemoglobin level remained elevated (4.4% vs. 0.8% prior to exposure) (44). 230

Based on these observations, prohibiting a methylene chloride worker from smoking on the job would not be sufficient to protect the worker who smokes after he leaves work from the additive exposures of CO from methylene chloride and tobacco smoke. 233
234
235

Other Chemical Agents 238

Other chemical agents found in tobacco or the combustion of tobacco products and also found in the workplace are: acetone, acrolein, aldehydes, arsenic, cadmium, formaldehyde, hydrogen sulfide, ketones, lead, methyl nitrite, nicotine, nitrogen dioxide, phenol, polycyclic compounds (26). 241
242
243

Smoking may contribute to an effect comparable to that resulting from exposure to toxic agents found in the workplace, thus causing an additive biological effect. 247
248
249

Coal Dust 252

Coal dust and cigarette smoking appear to act in an additive fashion to 255

1005058078

produce obstructive airway disease. Although dust exposure alone plays a significant role in the development of obstructive airway disease, there is a significantly higher prevalence of obstructive airway disease in smoking miners than in non-smoking miners with the same dust exposure (45). Flow volume curve data from the use of sophisticated pulmonary function techniques suggest that non-smoking miners with dust induced chronic obstructive airway disease have decreased flow rates at higher lung volumes, whereas, smoking miners have decreased flow rates at all lung volumes (46).

Cotton Dust

Many investigators have noted that among cotton workers, cigarette smokers show increased prevalence of byssinosis when compared to non-smoking cotton workers (47,48,49,50). Cotton dust inhalation produces an acute clinical syndrome consisting of chest tightness, cough, and shortness of breath in cotton workers (51). This was formerly known as "Monday fever" since symptoms develop on the first day of work after an absence. The clinical syndrome may be accompanied by significant reduction in pulmonary function (52). The acute clinical and functional abnormalities produced by cotton dust gradually become more frequent as the disease progresses, eventually resulting in chronic obstructive airways disease (51).

In the acute phase of the illness there is a significantly greater diminution in pulmonary function in smokers than in non-smokers (48) and

1005058079

the relationship of cotton dust and smoking to pulmonary dysfunction 280
appears to be additive.

In the more severe phase of chronic obstructive airway disease, the 282
relationship between smoking and cotton dust exposure appears to be 283
synergistic (48).

Beta Radiation 286

In studies in mice when both beta radiation and cigarette tar were 289
applied to produce carcinomas in the skin, cancers appeared six to seven 290
months earlier than when radiation was administered alone. The 291
shortened latent period gave an illusion of synergism which was reported 292
in a preliminary analysis based on tumor yield at 18 months. However, 293
at the conclusion of the experiment, the authors felt there was actually 294
nothing more than an additive biological effect of cigarette "tar" and 295
beta radiation (53).

Chlorine 298

Exposure to chlorine and cigarette smoke may cause an additive 301
biological effect. Chester et al. (54) examined 139 men in a plant 302
producing chlorine and sodium hydroxide by electrolysis of brine. 303

1005058080

Fifty-five of the 139 workers had been accidentally exposed one or more 304
times to chlorine at high concentrations and had required oxygen therapy 305
at least once during their employment. The maximal mid-expiratory flow 307
(MMF) values of workers with accidental chlorine exposure was compared 308
with those of non-exposed workers for smokers and non-smokers. A 309
significant difference in maximal mid-expiratory flow was seen when 310
chlorine and smoking were considered as additive toxic agents. Maximal 311
mid-expiratory flow values decrease from unexposed non-smokers (4.36) to 312
unexposed smokers (4.13) to exposed non-smokers (4.10) and to exposed 313
smokers (3.57).

Maximal Mid-Expiratory Flow Values of Workers 317
with Accidental Chlorine Exposure by Smoking 318
Category Compared to Non-exposed Workers. 319

	Exposed	Non-Exposed	320
			321
Smoker	3.57	4.13	322
			323
Non-Smoker	4.10	4.36	324
			325
			326
			327
			328

1005058081

Capodaglio et al. (55) studied the diffusing capacity of the lung in 331
workers employed in a plant for electrolytic production of chlorine and 332
soda. He compared 52 exposed workers to 27 unexposed workers. The 333
diffusing capacity of the lung was significantly lower in exposed
smokers than in non-exposed smokers ($P \leq 0.02$), lower in exposed smokers 334
than in exposed non-smokers, and lower in exposed smokers than in 335
unexposed non-smokers ($P \leq 0.03$). 336
These studies show the additive effects of cigarette smoking and 339
chlorine exposure.

Smoking may act synergistically with toxic agents found in the workplace 342
to cause a more profound effect than that resulting from smoking and the 343
agent added together. 344

ASBESTOS AND SMOKING 349

Asbestos provides one of the most dramatic examples of adverse health 351
effects resulting from interaction between the smoking of tobacco 352
products and an agent used in the workplace. Asbestos, the generic term 354
used to describe chain-silicates, was first used in Finland to 355

1005058082

strengthen clay pottery about 2500 B.C. (56). Modern industrial use of
asbestos is relatively more recent, dating from 1880 when it was used to
make heat and acid resistant fabrics (57,58). From that beginning its
usefulness has grown immensely with its output having increased over one
thousand-fold in the past sixty years (56).

With increasing industrial importance has come an increasing awareness
of the adverse health consequences incurred by working with asbestos.
Early in the twentieth century asbestosis was first reported and
subsequent individual observations and epidemiological studies have well
defined the association of this non-malignant respiratory disease and
asbestos exposure. In 1935 Lynch and Smith reported a suspected
association between asbestosis and lung cancer (59). Succeeding
epidemiologic studies have given significant support to these early
reports.

In 1968 a study of insulation workers by Selikoff et al. (60) defined
cigarette smoking as an additional hazard to the health of workers
exposed to asbestos. In a study of asbestos insulation workers,
Selikoff found that of 87 non-smokers, none died of bronchogenic
carcinoma, while 24 out of 283 cigarette smokers died of bronchogenic
carcinoma. This study suggested that asbestos workers who smoke have 8
times the lung cancer risk of all other smokers and 92 times the risk of
non-smokers not exposed to asbestos. This same group of insulation
workers were restudied five years later (61). At that time 41 of the

1005058083

283 smokers had died of bronchogenic cancer. In a larger study 379
involving 11,656 insulation workers in the United States and Canada, 134 380
deaths due to lung cancer were found among 9590 men with a history of 381
regular cigarette smoking. Of the 2066 non-cigarette smokers followed 382
over the same five year period, two deaths were due to lung cancer. 383

Over a ten year period, Berry et al. (62) studied 1300 male and 480 386
female asbestos factory workers in whom a smoking history was known. 387
The male and female groups were then evaluated considering whether they 388
had low to moderate or high asbestos exposure. The researchers found no 389
significant excess deaths from lung cancer in either smoking or non- 390
smoking groups at low to moderate exposures. However, a highly
significant increase in lung cancer deaths was seen in the severely 391
exposed who also smoked. 392

The above mentioned studies and other similar studies have shown that 395
cigarette smoking and asbestos exposure together are associated with
extremely high rates of lung cancer. But what role does each play in 396
this process? Two general hypotheses have been proposed to answer this 397
question (62). The additive hypothesis suggests that asbestos exposure 398
and cigarette smoking act independently to produce lung cancer and that 399
the excess risk seen when both are experienced together is due to the 400
sum of their risks. The multiplicative (synergistic) hypothesis 401
contends that each of the involved risk factors has a certain value for 402
its risk and that the product of these two risks (asbestos exposure x 403

1005058084

cigarette smoking) describes how they work together to bring about a
certain result (lung cancer). Selikoff's data suggests a synergistic
effect. However, in the study by Berry et al. (62), the male data does
not fit either hypothesis while the female data easily supports the
multiplicative hypothesis. A more recent study by Martischnig et al.
(63) of 201 men with confirmed bronchial carcinoma was much less
consistent with the multiplicative hypothesis and pointed more closely
to the additive hypothesis. Regardless of whether the action is
additive or synergistic, a substantial risk faces smokers who are exposed
to asbestos.

Other neoplasms have been associated with exposure to asbestos but
appear to be independent of smoking habits. Eighty-five to ninety
percent of mesothelioma have been attributed to exposure to asbestos
(64). The relationship of pleural and peritoneal mesothelioma to
smoking and asbestos exposure was investigated by Hammond and Selikoff
(61). Calculations from their studies reveal 0.38 deaths from pleural
mesothelioma per 1000 man years of observation among asbestos exposed
cigarette smokers and 0.39 for exposed non-smokers. Rates for
peritoneal mesothelioma were 0.73 for smokers and 0.83 for non-smokers
(65).

In 1971 Weiss (66) explored the relationship of asbestosis to cigarette
smoking. He examined 100 asbestos textile workers by chest x-ray and

1005058085

questionnaire. Pulmonary fibrosis was found in 40% of 75 workers who smoked and 24% of 25 non-smokers. Weiss determined that age, sex, and duration of exposure to asbestos were not responsible for the difference noted. Seventy-three of the above cigarette smokers were then questioned concerning amount and duration of smoking. The prevalence of fibrosis was 23% of 13 workers who smoked less than one pack per day and 43% of 60 who smoked one or more packs per day. Of 18 workers who smoked a pack or more per day for less than 20 years and had less than 20 years of asbestos exposure, 28% had fibrosis. Of 19 workers who smoked more than 20 years and with more than 20 years of exposure to asbestos, 74% had fibrosis. This study demonstrates that the prevalence of pulmonary fibrosis increases with increasing amount and duration of cigarette smoking and with increasing duration of exposure to asbestos. Due to the small size of the group he was working with, Weiss was unable to determine whether cigarette smoking and asbestos exposure were working in an additive or multiplicative manner. A study recently published by Weiss and Theodos indicates that type of asbestos as well as smoking habits are factors in the development of pleuropulmonary disease in asbestos workers (67).

In summary, workers exposed to tobacco smoke and asbestos experience far greater levels of lung cancer than would be expected from the contribution of either tobacco smoke or asbestos alone. However, other adverse health effects of occupational exposure to asbestos (e.g.,

1005058086

mesothelioma) appear to be independent of smoking habits. Thus, smoking 453
varies in its contribution to the development of different adverse 454
health effects resulting from occupational exposure to a particular 455
occupational agent.

Exposures in the Rubber Industry 460

In a study of rubber workers, Lednar et al. (68) reported that smokers 462
exposed to fumes and dust, particularly talc and carbon black, had a 463
significantly higher risk of developing a pulmonary disability than did 464
non-smokers. The combination of smoking and occupational exposure 466
significantly elevated the probability of developing an early pulmonary 467
disability. The authors reported that exposure to dust and smoking was 468
associated with 10 to 12 times the risk of pulmonary disability 469
retirement as a non-smoking, non-occupationally exposed rubber worker. 470
This elevated risk was found where there were exposures to respirable 471
particulates and/or solvents. This study suggests that smoking and 472
occupational exposures in the rubber industry are synergistic since the 473
authors report that a rubber worker who smoked and was exposed to talc 474
had an excess relative risk of 3.40 whereas an excess relative risk of 475
1.77 would be expected if the effects of smoking and work exposure were 476
additive. The mechanism of this interaction is not yet understood. 477

1005058087

Radon Daughters

479

A substantial excess of lung cancer, reduced pulmonary function, and emphysema has been reported among uranium miners (69). The excess has been attributed primarily to irradiation of the tracheobronchial epithelium by alpha particles emitted during the decay of radon and its daughter products. In a study of uranium miners, Archer et al. (70) found that respiratory cancer rates among smoking and non-smoking uranium miners were six to nine times greater than among non-miners with similar smoking habits. The lung cancer rate for nonsmoking uranium miners was 7.1 per 10,000 person years compared to 1.1 for non-miners who did not smoke. The lung cancer rate for uranium miners who smoked was 42.2 per 10,000 person years compared to 4.4 for non-miners who smoked 2 or more packs of cigarettes a day (Figure 1). There was also a definite association between the prevalence of emphysema and the cumulative amount of cigarettes smoked as well as with accumulative radiation exposure.

482

483

484

485

486

487

488

489

490

491

492

493

494

Smoking may contribute to accidents in the workplace

496

Studies have shown that smoking contributes to accidents in the workplace. In a nine-month study of job accidents, the total accident rate was more than twice as high among smokers as among non-smokers (71). Other authors have suggested that injuries attributable to

498


499

500

501

1005058088

Fig. 1 - Respiratory Cancer Rates Among Uranium Miners by Cigarette Usage and Radiation Exposure Compared with Rates Among Non-Miners*



*From:

Archer V.E., Wagoner, J.K., and Lunden, F.E., Jr.
 "Uranium Mining and Cigarette Smoking Effects on Man".
 Journal of Occupational Medicine, 15(3): 204-211, March 1973.

1005058089



Editor IRVING R. TABERSHAW, M.D.

Executive Editor DORIS L. FLOURNOY, M.S.

JOURNAL OF OCCUPATIONAL MEDICINE

OWNED AND PUBLISHED BY THE AMERICAN OCCUPATIONAL MEDICAL ASSOCIATION

150 NORTH WACKER DRIVE • CHICAGO, ILLINOIS 60606 • 312/782-2166


June 14, 1978

Harvey P. Stein, Ph.D.
Office of Extramural Coordination
and Special Projects
Dept. of Health, Education and Welfare
NIOSH
5600 Fishers Lane
Rockville, Maryland 20852

Dear Dr. Stein:

We are pleased to grant you permission to reproduce Figure 1 from the article "Uranium Mining and Cigarette Smoking Effects on Man" by Drs. Archer, Wagoner, and Lundin as outlined in your letter of June 8. This permission is contingent upon the authors' approval and is non-exclusive for one time use only with appropriate credit to the authors and the Journal of Occupational Medicine.

Sincerely yours,


Doris Flournoy
Executive Editor

DF:bp

1005058090



DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE
PUBLIC HEALTH SERVICE

HEALTH SERVICES AND MENTAL HEALTH ADMINISTRATION

Rm. 433, USPO & Courthouse Bldg., 350 So. Main
Salt Lake City, Utah 84101

NATIONAL INSTITUTE FOR OCCUPATIONAL
SAFETY AND HEALTH

July 5, 1978


Harvey P. Stein, Ph.D.
Senior Chemist
Technical Evaluation and
Review Branch
Office of Extramural
Coordination and Special Projects
DHEW, CDC, NIOSH
5600 Fishers Lane
Rockville, Maryland 20852

Dear Dr. Stein:

Thank you for sending me a copy of your letter to D. L. Flournoy
of J.O.M. requesting permission to reproduce a figure from my
article, "Uranium Mining and Cigarette Smoking Effects in Man".

Subject to approval by the editor of J.O.M., I am happy to grant
you my approval.

Sincerely,


Victor E. Archer, M.D.
Medical Director

1005058091

smoking were caused by loss of attention, preoccupation of the hand for 502
smoking, irritation of the eyes, and cough (72). 503

Smoking can also contribute to fire and explosions in occupational 506
settings where flammable and explosive chemical agents are used. In 508
many of these areas smoking is prohibited. For example, smoking is not 509
permitted in coal mines and miners are personally fined if in violation
of this provision.

EXAMPLES WHERE ACTION BETWEEN SMOKING AND OCCUPATIONAL EXPOSURE HAS BEEN 516
SUGGESTED OR ONLY HYPOTHESIZED

Cadmium 519

Several studies of the effects of occupational exposure to cadmium on 522
smokers and non-smokers have been conducted (73,74,75,76,77). Pulmonary 523
function is poorer in smokers than in non-smokers exposed to cadmium and 524
smokers also had a higher incidence of proteinuria than did non-smokers 525
in a cadmium exposed population in a Swedish battery factory. An 526
additive rather than a potentiating effect seems more likely from the 527
limited data.

1005058092

Chloromethyl Ether 530

A group of 129 men in a chemical plant where chloromethyl ether was used 533
were screened by 70 mm chest photofluorograms and questionnaires 534
regarding age, smoking habits, and respiratory symptoms at intervals 535
averaging 8.5 months for five years and follow-up for an additional five 536
years (78). Each job classification was ranked according to degree of
exposure to chloromethyl ether and an exposure index was calculated for 538
each man by cumulating the total exposure.

Chronic cough and expectoration showed a dose response relationship to 541
chemical exposure. Chronic cough was also related to smoking but for 542
each smoking category, chronic cough was more common for exposed than. 543
for unexposed men.

The 10 year incidence of lung cancer was dose related to chemical 546
exposure but not related to cigarette smoking. All cancers were small 547
cell carcinomas, occurred in men younger than 55 and had an induction- 548
latent period of 10 to 24 years. The 10 year mortality rate in this 549
group of workers was 2.7 times expected and lung cancer accounted for 550
the excess number of deaths.

Bronchogenic carcinomas linked to cigarette smoking are most often 553
squamous cell in type with long induction-latent periods and tend to 554
occur after the age of 60. The cancers which occur in workers exposed 555

1005058093

to chloromethyl ether are small cell in type, have short induction- 556
latent periods and tend to appear before the age of 55. The absence of 557
a relationship between cigarette smoking and lung cancer in this study
may be due to the competing effect of chloromethyl ether which results 559
in lung cancer in exposed workers before the long-term carcinogenic
effect of cigarette smoking could be demonstrated. However, cough 560
related to cigarette smoking appears earlier in exposed workers, thus 561
demonstrating the action of cigarette smoking with exposure to 562
chloromethyl ether in the development of chronic cough symptoms. This 563
case study also points up the complex issues involved in understanding 564
the actions between smoking and occupational exposures. 565

beta-Naphthylamine and other aromatic amines 568

Doll et al. found an excess risk of bladder cancer in a series of 570
studies (79,80) of men employed in coal gas production in England and 571
Wales. Most of the gas workers were smokers. Chemical studies showed 573
that inside the retort houses gas workers inhaled beta-naphthylamine and 574
other aromatic amines (known bladder carcinogens). Since aromatic 575
amines are also found in cigarette smoke (26), the gas workers who 576
smoked received exposure to bladder carcinogens from two sources. This 578
evidence is speculative but points out the need to assess the action 579
between smoking and exposure to aromatic amines.

1005058094

TRENDS IN SMOKING HABITS AND IN MORBIDITY AND MORTALITY RATES IN 583
OCCUPATIONAL GROUPS 584

Surveys (81) have shown male blue-collar workers are much more likely to 587
smoke cigarettes than white-collar workers. While only 37% of white- 589
collar workers were reported in 1973 to be current smokers, 51% of those
in blue-collar occupations then smoked. Also, more ex-smokers are found 590
among white-collar workers than among blue-collar workers (35% and 28% 591
respectively). Smoking among white-collar workers dropped from 48% to 592
37% between 1966 and 1970; during the same time period smoking among 593
blue-collar workers dropped from 62% to 51%.

The pattern among female employees is quite different (81). There was 595
little difference in smoking rates between white- and blue-collar female 596
workers, 36% and 38% respectively, in the 1973 report. In addition, the 597
smoking rates for 1966 were the same as those for 1970 in both groups of 598
female workers. During the period studied, the increased cessation of 599
smoking among female workers was offset by the increased initiation of 601
smoking in the same group. 602

In a study by Boucot et al. (82), one hundred twenty-one new lung 604
cancers developed among 6,136 men aged 45 and over who volunteered to 606
report semiannually for chest x-rays and answer questionnaires about
symptoms, smoking habits, etc., over a 10-year period beginning in 1951. 607
The risk of developing lung cancer increased with increasing age, was 608

1005058095

higher in non-whites than in whites, and bore a dose-response 609
relationship to cigarette smoking. The highest lung cancer risk was 611
among asbestos workers, 42.9/1000 man-years. The risk was 2.2/1000 man- 612
years for men in occupational categories not thought to be associated 613
with an increased risk of lung cancer. Occupational categories showing 614
somewhat increased risk were metal workers, cooks, and automobile 615
drivers. A higher percentage of whites, than non-whites (86.5% vs 616
77.4%), worked in occupations not thought to be at increased lung cancer 617
risk.

The smoking habits in various occupational groups demonstrate ample 619
opportunity for interaction between cigarette smoking and physical and 621
chemical agents in the workplace. In general, those who have the
highest smoking rates also have the highest risk for industrial 622
exposures. Both the consumption of tobacco products and exposure of 623
industrial agents increased steadily from 1920 to 1960. This is 624
reflected in certain mortality trends. For example the United States 625
age-adjusted mortality rate from carcinoma of the pancreas has been 626
reported to have risen from 2.9 to 8.2 per 100,000 population from 1920
to 1965, an increment of 283%. The rise was found to be real and 627
threefold in magnitude when adjustments were made for the aging of the 628
population. A literature review on pancreatic cancer was conducted by 629
Krain to determine real causes or associations for pancreatic cancer.
His report indicated that only the data on industrial carcinogen 631
exposure and cigarette smoking show both the trend and the statistical 632

1005058096

mangitude of association to consider them as real causes or
associations (83). 633

Since 1966 the consumption of tobacco products has decreased in blue 636
collar workers while the number of industrial exposures have continued
to increase (81,84). The increasingly higher rates of lung cancer in 637
non-white males, independent of smoking habits, may reflect the late 638
entry of non-whites into industrial settings and the fact that they have 639
jobs with higher risk for occupational exposure to toxic agents. 640

SUMMARY AND RECOMMENDATIONS 642

Although precise relationships between smoking and occupational 644
exposures cannot be quantified, the necessary data are beginning to 645
accumulate.

From 1920 to 1966 both tobacco consumption increased as did the 647
introduction into the workplace of chemicals with unknown biologic 648
effects. Workers with the greatest risk of exposure to industrial 649
agents also had the highest smoking rates. Since 1966 the consumption 650
of tobacco products has decreased in male blue collar workers while the 651
introduction of new chemicals into the workplace has continued to 652
increase.

1005058097

At least six different ways have been illustrated by which smoking may
act with physical and chemical agents in the workplace to produce
adverse health effects. These actions need not be mutually exclusive
and exposure to multiple physical and chemical agents in the workplace
can compound these various types of actions.

The examples of the interactions between the smoking of tobacco products
and industrial exposures cited in this report indicate that a
curtailment of smoking in certain occupational settings would contribute
to the reduction of specific disease processes. NIOSH has therefore
recommended in certain circumstances that workers exposed to particular
agents refrain from smoking. However, it is important to note that in
some situations (for example, radon daughters and chloromethyl ether),
the contribution of occupational exposures to adverse health effects was
greater than the contribution of cigarette smoking. Therefore, the
curtailment of smoking in the workplace should not be done in lieu of
curtailing occupational exposures to physical and chemical agents.

Recommendations

1. Studies on the health effects from smoking should take occupational
exposures into consideration and vice versa. Whenever possible,
studies should include data on occupationally exposed non-smoking
as well as unexposed smoking controls.

1005058098

2. The increasing rates of lung cancer in non-white males compared to white males should be investigated further with respect to occupational exposures and smoking habits. 680
681
3. The change in smoking habits of blue collar workers over the last decade provides an opportunity to more critically assess the contribution of smoking vs. occupational exposure to certain disease states. Prospective cohorts should be identified and followed for this purpose. 683
684
685
686
4. Workplace agents should be identified which interact with the smoking of tobacco to produce adverse health effects. 689
5. More studies on the modes of synergism between smoking and occupational exposures are needed. 692
6. The impact of the combination of smoking and workplace exposures upon reproductive disorders merits further study. 694
695
7. The impact of smoking in the workplace upon accidents merits further study. 697
698
8. The lack of information on the effect of side stream smoke in the development of occupational disease in non-smoking workers merits attention. 701

1005058099

REFERENCES

1. National Institute for Occupational Safety and Health. Criteria
for a Recommended Standard....Occupational Exposure to
Decomposition Products of Fluorocarbon Polymers. U.S.
Department of Health, Education, and Welfare, Public Health
Service, Center for Disease Control, National Institute for
Occupational Safety and Health, DHEW (NIOSH) Publication No. 77-
193, September 1976 123pp. National Technical Information
Service, PB274727.
2
3
4
5
7
8
9
10
11
12
13
2. William, N., Atkinson, G.W., Patchefsky, A.S. Polymer fume fever:
not so benign. Journal of Occupational Medicine 16(8): 519-522,
August 1974.
16
17
3. Evans, E.A. Pulmonary edema after inhalation of fumes from
polytetrafluoroethylene (PTFE). The Journal of Occupational
Medicine 15(7): 599-601, July 1973.
19
20
21
4. Robbins, J.J., Ware, R.L. Pulmonary edema from teflon fumes. The
New England Journal of Medicine 271(7): 360-361, August 13,
1964.
24

1005058100

5. Harris, D.K. Polymer fume fever. The Lancet 261(6992): 1008- 26
1011, December 1, 1951. 27
6. Touey, G.P., and Mumpower, R.C. Measurement of the combustion- 29
zone temperature of cigarettes. Tobacco 144(8): 18-22, February 30
22, 1957.
7. Bruton, D.M. Polymer fume fever. British Medical Journal 32
1(5542): 757, March 25, 1967. 33
8. Barnes, R., and Jones, A.T. Polymer-fume fever. Medical Journal 35
of Australia 2(2): 60-61, July 8, 1967. 36
9. Wegman, D.H., and Peters, J.M. Polymer fume fever and cigarette 38
smoking. Annals of Internal Medicine 81(1): 55-57, July 1974. 39
10. Adams, W.G.F. Polymer fume fever due to inhalation of fumes from 42
polytetrafluoroethylene. Transactions of the Association of 43
Industrial Medical Officers 13: 20-21, 1963.
11. Barlas, J.A. Polymer fume fever in a printing works.- British 45
Medical Journal 1(5539): 567, March 4, 1967. 46
12. Kuntz, W.D., McCord, C.P. Polymer-fume fever. Journal of 48
Occupational Medicine 16(7): 480-482, July 1974. 49

1005058101

13. Sherwood, R.J. The hazards of fluon. Transactions of the 51
Association of Industrial Medical Officers 5: 10-12, 1955. 52
14. Williams, N., and Smith, F.K. Polymer-fume fever. An elusive 54
diagnosis. Journal of the American Medical Association 219(12): 55
1587-1589, March 20, 1972.
15. Clayton, Jr., J.W. Fluorocarbon toxicity and biological action. 57
Fluorine Chemistry Reviews 1(2): 197-252, 1967. 58
16. National Institute for Occupational Safety and Health. Criteria 60
for a Recommended Standard....Occupational Exposure to 61
Formaldehyde. Department of Health, Education, and Welfare, 62
Public Health Service, Center for Disease Control, National 63
Institute for Occupational Safety and Health, DHEW (NIOSH) 64
Publication No. 77-126, December 1976, 177 pp. National 65
Technical Information Service, PB273805.
17. National Institute for Occupational Safety and Health. Criteria 67
for a Recommended Standard....Occupational Exposure to Boron 68
Trifluoride. U.S. Department of Health, Education, and Welfare, 69
Public Health Service, Center for Disease Control, National 70
Institute for Occupational Safety and Health, DHEW (NIOSH) 71
Publication No. 77-122, December 1976, 95 pp. National Technical 72
Information Service, PB274747.

1005058102

18.	National Institute for Occupational Safety and Health. Criteria	74
	for a Recommended Standard....Occupational Exposure to Organotin	75
	Compounds. U.S. Department of Health, Education, and Welfare,	76
	Public Health Service, Center for Disease Control, National	77
	Institute for Occupational Safety and Health, DHEW (NIOSH)	78
	Publication No. 77-115, November 1976. 196 pp. National	
	Technical Information Service, PB223113.	79
19.	National Institute for Occupational Safety and Health. Criteria	81
	for a Recommended Standard....Occupational Exposure to Methyl	82
	Parathion. U.S. Department of Health, Education, and Welfare,	83
	Public Health Service, Center for Disease Control, National	84
	Institute for Occupational Safety and Health, DHEW (NIOSH)	
	Publication No. 77-106, September 1976, 189 pp. National	85
	Technical Information Service, PB274191.	
20.	National Institute for Occupational Safety and Health. Criteria	87
	for a Recommended Standard....Occupational Exposure to Dinitro-	88
	Ortho-Cresol. U.S. Department of Health, Education, and Welfare,	89
	Public Health Service, Center for Disease Control, National	90
	Institute for Occupational Safety and Health, DHEW (NIOSH)	
	Publication No. 78-131, February 1978, 157 pp.	91

1005058103

21. National Institute for Occupational Safety and Health. Criteria 93
for a Recommended Standard....Occupational Exposure to Carbaryl. 94
U.S. Department of Health, Education and Welfare, Public Health 95
Service, Center for Disease Control, National Institute for 96
Occupational Safety and Health, DHEW (NIOSH) Publication No. 77- 97
107, September 1976, 204 pp. National Technical Information
Service, PB273801.
22. National Institute for Occupational Safety and Health. Criteria 99
for a Recommended Standard....Occupational Exposure to Inorganic 100
Fluorides. U.S. Department of Health, Education, and Welfare, 101
Public Health Service, Center for Disease Control, National 102
Institute for Occupational Safety and Health, DHEW (NIOSH)
Publication No. 76-103, 1975. 191 pp. National Technical 103
Information Service, PB246692.
23. National Institute for Occupational Safety and Health. Criteria 105
for a Recommended Standard....Occupational Exposure to Inorganic 106
Mercury. U.S. Department of Health, Education, and Welfare, 107
Public Health Service, Health Services and Mental Health 108
Administration, National Institute for Occupational Safety and
Health, DHEW (NIOSH) Publication No. 73-11024, 1973. 138 pp. 109
National Technical Information Service, PB222223.

1005058104

24. Tola, S., Nordman, C.H. Smoking and blood concentrations in lead- 111
exposed workers and an unexposed population. Environmental 112
Research 13: 250-255, 1977. 113
25. Zielhuis, R.L., Stuik, E.J., Herber, R.F.M., Salle, H.J.A., 115
Verberk, M.M., Posma, F.D., Jager, J.H. Smoking habits and 116
levels of lead and cadmium in blood in urban women. 117
International Archives of Occupational and Environmental Health 118
39: 53-58, 1977.
26. U.S. Public Health Service. Smoking and Health - Report to the 120
Advisory Committee to the Surgeon General of the Public Health 121
Service, U.S. Department of Health, Education, and Welfare, 122
Public Health Service, Publication No. 1103.
27. Pettigrew, A.R., Fell, G.S. Simplified colorimetric determination 124
of thiocyanate in biological fluids, and its application to 125
investigation of the toxic amblyopias. Clinical Chemistry, 126
18(9): 996-1000, September 1972.
28. Radojicic, B. Oredivanje Rodanida U Mokraci U Radnika Izlozenih 128
Cijanidima (Determining thiocyanate in urine of workers exposed 129
to cyanides.) Arhiv Za Higijenu Rada I Toksikologiju 24: 227-
232, 1973.

1005058105

29. Fassett, D.W. Cyanides and nitriles. IN: Patty, F.A. Industrial 132
Hygiene and Toxicology, Second Revised Edition, Volume II. New 133
York, John Wiley and Sons, Inc., 1963, pp. 1991-2036.
30. Binkley, F. On the nature of serine dehydrase and cysteine 135
desulfurase. Journal of Biological Chemistry 150(1): 261-262, 136
September 1943.
31. Altschul, A.M., Abrams, R., Hogness, T.R. Cytochrome c 138
peroxidase. Journal of Biological Chemistry 136(3): 777-794, 139
December 1940.
32. Wood, J.L. Biochemistry. IN: Wood, J.L. Chemistry and 141
Biochemistry of Thiocyanic Acid and its Derivatives, New York, 142
Academic Press Inc., 1975, pp. 156-221. 143
33. National Institute for Occupational Safety and Health. Criteria 145
for a Recommended Standard....Occupational Exposure to Hydrogen 146
Cyanide and Cyanide Salts. U.S. Department of Health, 147
Education, and Welfare, Public Health Service, Center for 148
Disease Control, National Institute for Occupational Safety and 149
Health, DHEW (NIOSH) Publication NO. 77-108, October 1976, 204 150
pp. National Technical Information Service, PB266230. 151

1005058106

34. Chevalier, R.B., Krumholz, R.A., and Ross, J.C. Reaction of non- 154
smokers to carbon monoxide inhalation Cardiopulmonary responses 155
at rest and during exercise. Journal of the American Medical 156
Association 198(10): 1061-1064, December 5, 1966.
35. Judd, H.J. Levels of carbon monoxide recorded on aircraft flight 158
decks. Aerospace Medicine 42(3): 344-348, March 1971. 159
36. Osborne, J.S., Adamek, S., and Hobbs, M.E.: Some Components of 161
the Gas Phase of Cigarette Smoke. Annals of Chemistry, Vol. 28, 162
pp 211-215, 1957. 163
37. Goldsmith, J.R., Terzaghi, J., and Hackney, J.D. Evaluation of 165
fluctuating carbon monoxide exposures. Archives of 166
Environmental Health 7(6): 647-663, December 1963. 167
38. Jones, J.G., Walters, D.H. A study of carboxyhemoglobin levels in 170
employees at an integrated steelworks. Annals of Occupational
Hygiene 5: 221-230, 1962 171
39. Knelson, J.H.: Cardiovascular Effects During Low Level CO 173
Exposure, Paper presented to the Committee on Motor Vehicle 174
Emissions, NAS-NRC, Washington, D.C., February 10, 1972. 175

1005058107

40. Horvat, M., Yoshida, S., Prakash, R., Marcus, H.S., Swan, H.J.C., 177
Ganz, W. Effect of oxygen breathing or pacing-induced angina 178
pectoris and other manifestations of coronary insufficiency. 179
Circulation 45(4): 837-844, April 1972. 180
41. Hosey, A.D. Priorities in developing criteria for "breathing air" 183
standards. Journal of Occupational Medicine 12(2): 43-46, 1976
42. Fodor, G.G., Prajsnar, D., Schlipkoter, H.W. Endogene Co-Bildung 186
durch inkorporierte Halogenkohlenwasserstoffe der Methanreihe
(Endogenous CO Formation by Encorporated Halogenated 187
Hydrocarbons of the Methane Series.) Staub Reinhaltung der 188
Luft, (6): 258-259, June 1973.
43. Astrand, I., Ovrum, P., Carlsson, A. Exposure to methylene 190
chloride. I. Its concentration in alveolar air and blood during 192
rest and exercise and its metabolism. Scandinavian Journal of
Work, Environmental and Health 1(2): 78-94, June 1975. 193
44. Stewart, R.D., Hake, C.L. Paint-remover hazard. Journal of the 195
American Medical Association 235(4): 398-401, January 26, 1976. 196

1005058108

45. Kibelstis, J.A., Morgan, E.J., Reger, R., Lapp, N.L., Saaton, A., 200
Morgan, W.K.C. Prevalence of bronchitis and airway obstruction 201
in American bituminous coal miners. American Review of 202
Respiratory Disease 108(4): 886-893, 1973. 203
46. Hankinson, J.L., Reger, R.B., Morgan, W.K.C. Maximal expiratory 206
flows in coal miners. American Review of Respiratory Disease 207
116(2): 175-180, 1977.
47. Merchant, J.A., Lumsden, J.C., Kilburn, K. H., O'Fallon W.M., 209
Ujda, J.R., Germino, V.H., Hamilton, J.D. Dose response 210
studies in cotton textile workers. Journal of Occupational 211
Medicine 15(3): 222-230, March 1973. 212
48. Merchant, J.A., Lumsden, J.C., Kilburn, K.H., O'Fallon, W.M., 214
Ujda, J.R., Germino, V.H., Hamilton, J.D. An industrial study 216
of the biological effects of cotton dust and cigarette smoke 217
exposure. Journal of Occupational Medicine 15(3): 212-221,
March 1973.
49. Berry, G., McKerrow, C.B., Molyneux, M. K. B., Rossiter, C.E., 219
Tomblason, J.B.L. A study of the acute and chronic changes in 221
ventilatory capacity of workers in Lancashire cotton mills. -
British Journal of Industrial Medicine 30(1): 25-36, January 222
1973.

1005058109

50. Merchant, J.A., Kilburn, K.H., O'Fallon, W.M., Hamilton, J.D., 224
Lumsden, J.C. Byssinosis and chronic bronchitis among cotton 226
textile workers. *Annals of Internal Medicine* 76(3): 423-433,
March 1972. 227
51. Harris, T.R., Merchant, J.A., Kilburn, K.H., Hamilton, J.D. 229
Byssinosis and respiratory diseases of cotton mill workers. 230
Journal of Occupational Medicine 14(3): 199-206, March 1972. 231
52. Merchant, J.A., Halprin, G.M., Hudson, A.R., Kilburn, K.H., 233
McKenzie, W.N., Hurst, D.J., Bernazohn, P. Responses to cotton 234
dust. *Archives of Environmental Health* 30(5): 222-229, May 235
1975.
53. Cowdry, E.V., Croninger, A., Solaric, S., Suntzeff, V. Combined 237
action of cigarette tar and beta radiation in mice. *Cancer* 238
14(2): 344-352, March-April 1961. **1005058110**
54. Chester, E.H., Gillespie, D.C., Krause, F.D. The prevalence of 240
chronic obstructive pulmonary disease in chlorine gas workers. 241
American Review of Respiratory Disease 99(3): 365-73, 1969. 242
55. Capodaglio, E., Pezzagno, G., Bobbio, G.C., Cazzoli, F. Indagine 244
Sulla Funzionalità Respiratoria Di Lavoratori Addetti A 245
Produzione Elettrolitica Di Cloro E Soda. (Investigation of 246
the respiratory function of workers employed in electrolytic 247
production of chlorine and soda.) *Medicina Del Lavoro* 60(3): 248
192-202, 1969.

56. Stell, P.M., McGill, T. Exposure to asbestos and laryngeal carcinoma. *Journal of Laryngology and Otology* 89(5): 513-517, 1975. 251 252
57. Hendry, N.W. The geology, occurrences, and major uses of asbestos. *Annals of The New York Academy of Sciences*, 132(1): 12-22, December 31, 1965. 254 255
58. Hueper, W.C. Occupational and Environmental Cancers of the Respiratory System, Recent Results in Cancer Research 3. Berlin, Springer-Verlag, 1966, p. 38-56. 257 259
59. Lynch, K.M., Smith, W.A. Pulmonary asbestosis III: Carcinoma of the lung in asbesto-silicosis. *American Journal of Cancer* 24(1): 56-64, May 1935. 262
60. Selikoff, I.J., Hammond, E.C., Churg, J. Asbestos exposure, smoking, and neoplasia. *Journal of the American Medical Association* 204(2): 104-110, April 8, 1963. 264 265 266
61. Hammond, E.C., Selikoff, I.J. Relation of cigarette smoking to the risk of death of asbestos-associated disease among insulation workers in the United States. *International Agency for Research on Cancer Publication No.8*, 312-317, 1973. 269 270 271

1005058111

62. Berry, G., Newhouse, M.C., Turok, M. Combined effect of asbestos 274
exposure and smoking on mortality from lung cancer in factory
workers. The Lancet 2(7775): 476-479, 1972. 275
63. Martischning, K.M., Newell, D.J., Barnsley, W.C., Cowan, W.K., 277
Feinmann, E.L., Oliver, E. Unsuspected exposure to asbestos and 279
bronchogenic carcinoma. British Medical Journal 1(6063): 746-
749, March 19, 1977.
64. Wagner, J.C., Gilson, J.C., Berry, G., Timbrell, V. Epidemiology 282
of asbestos cancers. British Medical Bulletin 27(1): 71-76,
1971.
65. Selikoff, I.J. Cancer risk of asbestos exposure. IN: Hiatt, 286
H.H., Watson, J.D., Winston, J.A. Origins of Human Cancer. New 287
York, Cold Springs Harbor Laboratory, 1977, pp. 1765-1783.
66. Weiss, W. Cigarette smoking, asbestos, and pulmonary fibrosis. 290
American Review of Respiratory Disease 104(2): 223-227, August 291
1971
67. Weiss, W., Theodos, P.A. Pneuropulmonary disease among asbestos 293
workers in relation to smoking and type of exposure. British 294
Journal of Industrial Medicine 20(5): 341-345, May 1978. 295

1005058112

68. Lednar, W.M., Tyroler, H.A., McMichael, A.J., Shy, C.M. The 298
occupational determinants of chronic disabling pulmonary disease 299
in rubber workers. Journal of Occupational Medicine 19(4): 263- 300
268, April 1977.
69. Lunden, F.E., Wagoner, J.K., Archer, O.E.,: Radon Daughter 302
Exposure and Respiratory Cancer. Quantitative and Temporal 303
Aspects, NIOSH and NIEHS Joint Monograph No. 1, National 304
Technical Information Service, Springfield, Va. 1971. 305
70. Archer, V.E., Wagoner, J.K., and Lunein, F.E. Uranium mining and 307
cigarette smoking effects on man. Journal of Occupational 308
Medicine 15(3): 204-211, March 1972. 309
71. Yuste, P.C., Guevara, M.L. Influenza Del Fumar En Los Accidentes 311
Laborales. Encuesta Estadística. (The influence of Smoking on 312
industrial accidents.) Medicina y Seguridad Del Trabajo 21(84): 313
38-46, October-December 1973. 314
72. Naus, A., Engler, V., Hetyohova, M., Vavreckova, D. Work inquires 316
and smoking. Industrial Medicine and Surgery 35(10): 880-881, 317
October 1966.
73. Pescator, M., Adamson, E., Elender, C.G., Peterson, B. Steninger, 320
P.: Studies on Cadmium exposed women. Paper presented at the 321
XVIIIth International Congress on Occupational Health, Brighton, 322
England 1975.

1005058113

74. Kjelstrom, T., Ervin, P.E., Rahnster, B. Dose-response analysis 325
of cadmium induced tubular proteinuria. A study of urinary B - 326
muroglobulin excretion among workers in a battery factory. 327
Environmental Research 13(2): 303-317, April 1977. 328
75. Lauwerys, R.R., Buchet, J.P., Roels, H.A., Brouwers, J., Stanescu, 331
D. Epidemiological survey of workers exposed to cadmium. 332
Effect on lung, kidney and several biological indices. Archives 334
of Environmental Health 28(3): 145-48, March 1974.
76. Lauwerys, R. Buchet, J.P., Roels, H.: A Study of Pulmonary and 337
renal function in workers exposed to cadmium, Paper presented at 338
the XVIIIth International Congress on Occupational Health, 339
Brighton, England 1975.
77. Materne, D., Lauwerys, R., Buchet, J.P., Roels, H., Brouwers, J., 342
Stanesca, D. Investigations Sur Les Risques Resultant De 343
L'exposition Au Cadmium Dans Deux Entreprises D'utilisation Du 344
Cadmium. (Investigations on the risk resulting from exposure to 345
cadmium in two production plants and two plants using cadmium.) 346
Cahiers De Medecine Du Travail 12(1 and 2): 3-76, March-June 347
1975.

1005058114

78. Weiss, W., Chloromethyl ethers, cigarettes, cough and cancer. 351
Journal of Occupational Medicine 18 (3): 194-199, March 1976.
79. Doll, R., Vessey, M.P., Beasley, B.W.R., Buckley, A.R., Fear, 354
E.C., Fisher, R.E.W., Gammon, E.J., Gunn, W., Hughes, G.O., Lee, 355
K. and Smith, B.N. Mortality of gas workers. Final report of a 356
prospective study. British Journal of Industrial Medicine 357
29(4): 394-406, 1972. 358
80. Doll, R., Fisher, R.E.W., Gammon, E.J., Gunn, W., Hughes, G.O., 360
Tyrer, F.H., Wilson, W. Mortality of gasworkers with special 361
reference to cancers of the lung and bladder, chronic 362
bronchitis, and pneumoconiosis. British Journal of Industrial 363
Medicine 22(1): 1-12, 1965.
81. National Clearinghouse for Smoking and Health. Adult Use of 366
Tobacco 1970. U.S. Department of Health, Education, and 367
Welfare, Public Health Service, Center for Disease Control, 368
National Clearinghouse for Smoking and Health, June 1973.
82. Boucot, K.R., Weiss, W., Seidman, H., Carnahan, W.J., Cooper, D.A. 370
Philadelphia Pulmonary neoplasm research project. Basic risk 371
factors of lung cancer in older men. American Journal of 372
Epidemiology 95(1): 4-16, 1972.

1005058115

83. Krain, L.S. The rising incidence of carcinoma of the pancreas - 375
real or apparent? Journal of Surgical Oncology 2(2): 115-124, 376
1970.
84. Bridbord, K., French, J.G., Finklea, J. Toxic materials - 379
respiratory pathways. Proceedings International Conference on 380
Cybernetics and Society The Institute of Electrical and
Electronics Engineers, Inc. Catalog No: 77CH1259-1 SMC: 80-81, 381
1977.

1005058116